

**Guidelines For Prophylaxis For RSV Infections
In High-Risk Infants In Hawaii
(Developed by Consensus Committee – July 15, 2008)**

Patient Population

1. All children younger than 2 years old at the beginning of the season (see below for definition) with Chronic Lung Disease and Congenital Heart Disease requiring treatment/medical management within 6 months of the anticipated season (Born on or after September 15, 2006; continuing medical treatment after March 15, 2008).
2. All children born prematurely at 28 weeks gestation or earlier, who are less than 1 year chronological age at the beginning of the season (Born on or after September 15, 2007) .
3. All children born prematurely between 29 and 32 weeks gestation, who are less than 6 months chronological age at the beginning of the season (Born on or after March 15, 2008).
4. All children born prematurely between 33 and 35 weeks gestation requiring significant respiratory support in the neonatal period (positive pressure support) and having an additional risk factor (child care attendance, school-aged siblings, congenital abnormalities of the airways or severe neuromuscular disease), who are less than 6 months chronological age at the beginning of the season (Born on or after March 15, 2008)
5. There are several children with other illnesses in the Pediatric age group who may be considered for prophylaxis, although these are not FDA-approved indications for the use of these drugs (Respigam® and Synagis ®). The Committee recommends that Pediatricians evaluate these children on a case-by-case basis and, if necessary, in consultation with an appropriate Sub specialist.
6. All children after cardiopulmonary bypass and with indication for use of Synagis should be considered for additional prophylaxis after discharge. Further, children with cardiac disease undergoing cardiopulmonary bypass during the season and currently receiving prophylaxis should receive an additional dose of prophylaxis, within a few days after bypass. They should continue to receive subsequent prophylaxis until the end of the season.

Season

RSV infections occur all year round in our community. However, based on available epidemiological data, the incidence is significantly higher from September through February. For infants eligible for immunoprophylaxis, the beginning of the season will be considered **September 15, 2008**, and immunoprophylaxis should be continued to provide immunity through February 2009. The Committee will plan on meeting again in the last two weeks of January 2009 to evaluate the end date for the season.

Prophylaxis

1. Prophylaxis for infants identified by criteria reflected under patient population should be started between September 15, 2008 and September 30, 2008. Prophylaxis for children who qualify based on age at the beginning of the season should be continued for the duration of that season.
2. Prophylaxis should be continued to provide immunity at least until the end of February 2009.
3. Should a child develop RSV during the course of the season, prophylaxis should be continued after recovery until the end of the season.
4. The maximum number of doses should be limited to five unless the season is extended by the Consensus Committee.
5. The interval between the first and second dose should be as close as possible to 28 days. All subsequent dose intervals should be as close to 30 days as possible (range 28 – 35 days).

These recommendations are meant to be guidelines.

Additional factors that need to be considered are:

1. Education of the family that although Prophylaxis is not 100% effective, it may lead to a decrease in severity of subsequent illness. To this effect, consideration should be given to obtaining an informed consent prior to drug administration.
2. Family education with respect to:
 - a. Use of good hand-washing practices during the wet and cooler months
 - b. Avoiding exposure to smoke and dust especially passive smoke exposure in the presence of smokers in the family
 - c. Avoiding contact with ill persons especially those with respiratory symptoms
 - d. Avoiding unnecessary exposure to crowds
3. **This vulnerable population should also be considered for Influenza immunization in addition to RSV prophylaxis, if they are over the age of six months.**

The Committee welcomes comments from community pediatricians and other health care providers regarding RSV infections in their practices and the impact of these guidelines on the same. Communication with the Committee may be directed to any of the Committee members listed below, or by contacting:

Venkataraman Balaraman, MB, BS.
Department of Pediatrics
Kapi'olani Medical Center for Women and Children
1319 Punahou Street, Room 750
Honolulu, Hawaii 96826
Email: vbaraman@kapiolani.org

Committee Members

Raul Rudoy, M.D., Chair
Vernon Azuma, M.D.
Venkataraman Balaraman, MB, BS.
Marta Derieg, M.D.
James Griffith, M.D.
Matthew Ho, M.D.
Douglas Kwock, M.D.
Sarah Lentz-Kapua, M.D.
Connie Lin, M.D.
James Lin, M.D.
Wallace Matthews, M.D.
Frank Smith, M.D.
Charles Neal, M.D.
Brenda Nishikawa, M.D.
James Sim, M.D.
Rupert Chang, M.D.
Brian Wilson, M.D.
Robert Wotring II, M.D.